

EFFECT OF STRUCTURED PHYSICAL ACTIVITY PROGRAMME ON GROSS MOTOR FUNCTION OF ADOLESCENCE WITH CEREBRAL PALSY

CEBY GEORGE¹, T.I. MANOJ²

¹SPSS, MG University, Kottayam, INDIA.

²Kerala Agricultural University, Vellanikkara, Thrissur, Kerala, INDIA.

Email: manojti@gmail.com

How to cite this article: George, C., & Manoj, T.I. (December, 2015). Effect of structured physical activity programme on gross motor function of adolescence with cerebral palsy. Journal of Physical Education Research, Volume 2, Issue IV, 38-47.

Received: October 04, 2015

Accepted: December 12, 2015

ABSTRACT

The purpose of the study was to find the effect of a carefully structured, moderate physical activity program on gross motor function of children with cerebral palsy. To achieve this purpose 16 (all male only) children with cerebral palsy between age group of 13-18 years with GMFCS levels I and II were selected as subjects for the study. They were randomly assigned to control and experimental groups of eight each. A structured moderate physical activity program was implemented to control group thrice in a week for 16 weeks. The participants of control and experimental groups were administered with GMFM 88 by the scholars before the start of training program and after implementation of the program. The MANOVA results shows that, there was a statistically significant main effect in gross motor function measure dimensions based on control group performance, $F(1, 16) = 609.21, p < 0.000$; Wilk's $\Lambda = 0.002$, partial $\eta^2 = 0.998$. It can be conclude that, 16 week structured moderate physical activity program can make significant improvement in GMFM dimensions: lying and rolling, sitting and walking, running and jumping of adolescent male with cerebral palsy.

Keywords: Physical activity, gross motor function, exercise, cerebral palsy.

1. INTRODUCTION

Cerebral palsy (CP) describes a group of disorders of the development of movement and posture, causing activity limitation, that are attributed to non-progressive disturbances that occurred in the developing fetal or infant brain. The motor disorders of CP are often accompanied by disturbances of sensation,

Correspondence: T.I. Manoj, Ph.D., Dy. DSW, Kerala Agricultural University, Vellanikkara, Thrissur, Kerala, INDIA, Tel: +91 9447063339, Email: manojti@gmail.com

cognition, communication, perception, and/or behavior, and/or by a seizure disorder (Bax, Goldstein, Rosenbaum, Leviton, Paneth, Dan, ..., & Executive Committee for the Definition of Cerebral Palsy, 2005). Because of the impairments, many children and adolescents with CP have at least difficulty with activities such as walking independently, negotiating stairs, running, or navigating safely over uneven terrain (Styer-Acevedo, 1999). Improving one's ability to walk or to perform other functional activities are often the primary therapeutic goals for children with CP (Shepherd, 1995).

In India, services to the Cerebral Palsy (C.P.) population, historically has depended on the initiatives of parents, mostly hospital based which were evidently at best partial fragmentary and grossly inadequate. The Spastics Society of India was founded in 1972 at a time when very little was known about the complicated disorder of cerebral palsy. Initially it provided education and treatment services gradually broadening its scope to teacher training, vocational training of young adults, advocacy and awareness, support for parents and other professionals. Integrated Education of Disabled Children as a scheme was launched in 1979 with the objective of providing educational opportunity to Special Needs Children in regular schools, to facilitate their retention in the school system and to place children from special schools in common schools. The planning of IED may include extracurricular activities like drawing, painting, dance, music, sports, craft and indoor games. In fact, signs and symptoms of cerebral palsy, such as spastic muscles, poor balance and gross motor delays, will get worse without treatment/exercise. Physical activity program is concerned, no definite or integrated program was planned. To cope with the social necessity, children with CP needs appropriate public education to take place in the 'Least Restrictive Environment'. Child needs to be experienced with a highly individualized educational plan (Karande & Kulkarni, 2008). The process of developing an individualized educational plan for a child with CP begins with an assessment of the child's needs. After the assessment as per the guidelines the school has to work with the parent and others involved in the child's education and treatment to develop individualized educational plan. This plan must develop with the cooperative effort of a team of educational experts consisting professionals, regular teachers, special education teachers, different therapists, counselors, psychologists, experts in developmental screening and experts in teaching children with special needs. Parents also should be given enough opportunity to participate in the planning, execution and evaluation of the individualized educational plan (Paleeri, 2010).

Structured physical activity programs are those where the activity is conducted either in a group or individualized program format, under the supervision of a trained exercise leader or therapist. Structured physical activity is usually led by a qualified professional and adults, including sports

organizations, guide play, school sports programs. The program is to improve or maintain the fitness level of the individual by keeping the intensity above of the daily activity of the reference to a structured action plan related to repeated muscle movement occurs and energy consumption. Exercise in children with CP has often been avoided because of the concern about the negative effect of such effort on muscle spasticity and children's movement patterns (Bobath, 1971). Several factors have contributed to a recent shift in perspective about the use of exercise in children with CP. Studies evaluating the effect of exercise on children with CP reported no adverse effect on patterns of movement, flexibility, or spasticity (Verschuren, Ketelaar, Gorter, Helders, Uiterwaal, & Takken, 2007). These findings have influenced current practice.

Most exercise programs for children with CP are primarily designed for the lower extremity. The most common functions of the lower extremity tend to be gross motor activities that involve repetitive, reciprocal, coordinated motions of both extremities to move through space and that often require little conscious effort once under way (Damiano, 2006). There has been an increased interest in developing and implementing exercise programs that improve the cardiovascular fitness (aerobic and anaerobic capacity) and/or lower-extremity muscle strength of children with CP. A structured physical activity program systematically integrates the aerobic and anaerobic activities. The purpose of the study was to find the effect of a carefully structured, moderate physical activity program on gross motor function of children with cerebral palsy.

2. METHODS AND MATERIALS

2.1 Sample

Participants were selected from Spastics Society of Karnataka, Centre for Special Education, Bangalore. To achieve this purpose 16 (all male) children with cerebral palsy between age group of 13-18 years with GMFCS levels I and II were selected as subjects for the study. They were randomly assigned to control and experimental groups of eight each.

2.2 Instrumentation

The Gross Motor Function Measure (GMFM) developed by Russell, Rosenbaum, Wright, and Avery, (2013) is a standardized observational instrument designed and validated to measure change in gross motor function over time in children with cerebral palsy. GMFCS level is a rating of severity of motor function. The scoring key is meant to be a general guideline. However, most of the items have specific descriptors for each score. It is imperative that the guidelines contained

in the manual be used for scoring each item. The expanded GMFCS (2007) includes an age band for youth 12 to 18 years of age and emphasizes the concepts inherent in the World Health Organization's International Classification of Functioning, Disability and Health (ICF). General headings for each level includes: Level I - Walks without Limitations, Level II - Walks with Limitations, Level III - Walks Using a Hand-Held Mobility Device, Level IV - Self-Mobility with Limitations; May Use Powered Mobility, Level V - Transported in a Manual Wheelchair. GMFM 88 has five dimensions as follows: A - Lying and rolling, B- Sitting, C- Crawling and kneeling, D- Standing and E- Walking, running and jumping. Between 12th and 18th birthday, Level I: Youth walk at home, school, outdoors, and in the community. Youth are able to walk up and down curbs without physical assistance and stairs without the use of a railing. Youth perform gross motor skills such as running and jumping but speed, balance, and coordination are limited. Youth may participate in physical activities and sports depending on personal choices and environmental factors. Level II: Youth walk in most settings. Environmental factors (such as uneven terrain, inclines, long distances, time demands, weather, and peer acceptability) and personal preference influence mobility choices. At school or work, youth may walk using a handheld mobility device for safety. Outdoors and in the community, youth may use wheeled mobility when travelling long distances. Youth walk up and down stairs holding a railing or with physical assistance if there is no railing. Limitations in performance of gross motor skills may necessitate adaptations to enable participation in physical activities and sports. These dimensions will measure on four point system ranging from 0-3 was executed.

2.3 Procedure

A structured moderate physical activity program was implemented to control group thrice in a week for 16 weeks. Each session lasted for 50-60 minutes. The participants of control and experimental groups were administered with GMFM 88 by the scholars before the start of training program and after implementation of the program.

2.4 Statistical Analysis

For the analysis of the data descriptive as well as MANCOVA was used. All statistical function was performed with the help of Statistical Package of Social Sciences (SPSS) version 19 software.

3. RESULTS

The important characteristics of the data on pre-test and post-test on different dimensions of control and experimental groups are presented below in Table 1.

Table 1: Descriptive statistics of post test scores of dependent variables (GMFM dimensions)

Dimensions	Group	Mean	SD	N
Lying & Rolling	Control	20.63	6.36	8
	Experimental	30.38	8.05	8
Sitting	Control	26.25	8.82	8
	Experimental	36.75	6.86	8
Crawling & Kneeling	Control	16.50	7.21	8
	Experimental	21.63	6.25	8
Standing	Control	17.75	3.24	8
	Experimental	22.13	5.86	8
Walking, Running & Jogging	Control	26.00	5.70	8
	Experimental	40.37	7.26	8
Total GMFM	Control	21.43	5.63	8
	Experimental	30.25	6.44	8

Table 2: ANCOVA for Adjusted Posttest Score on Dependent variables

Dimensions	Type III Sum of Squares	df	Mean Square	F	Sig. (p-value)	Partial Eta Squared
Lying & Rolling- A	10.934	1	10.93	15.96*	0.003	0.639
Sitting- B	38.819	1	38.81	59.92*	0.000	0.869
Crawling & Kneeling- C	.006	1	0.01	0.01	0.920	0.001
Standing-D	1.007	1	1.01	0.61	0.455	0.063
Walking, Running & Jogging-E	3.889	1	3.88	5.18*	0.049	0.365
Total GMFM	6.187	1	6.18	12.31*	0.007	0.578

* Significant at 0.05 levels

The MANOVA results shows that, there was a statistically significant main effect in Gross Motor Function Measure Dimensions based on control group performance, $F(1, 16) = 609.213$, $p < .000$; Wilk's $\Lambda = 0.002$, partial $\eta^2 = .998$.

Analyses of Covariance (ANCOVA) were carried out for the posttest scores of five dimensions and Total GMFM scores separately to find out any differences in the adjusted posttest means scores on selected dimensions between control and experimental groups. The results on table 2 shows that, An ANCOVA showed statistically significant treatment effects on GMFM dimensions; Lying and rolling ($F(1, 16) = 5.96, p = .003$), Sitting ($F(1, 16) = 59.92, p = .000$), Walking, Running & Jogging ($F(1, 16) = 59.92, p = .000$) and Total GMFM ($F(1, 16) = 12.31, p = .007$) at the same time GMFM dimensions Crawling & Kneeling ($F(1, 16) = 0.011, p = .920$) and Standing ($F(1, 16) = 0.610, p = .455$) were found not statistically significant. Moreover, as per the Cohen effect size (Cohen, 1992) the dimensions A and B's Partial Eta Squared is larger than 0.40, which means that, the treatment made 63.9% and 86.9% variability in post-test scores.

Table 3: Pair-wise comparison of dependent variables adjusted post test means between control and experimental groups

Dependent Variable Group		Mean Difference	Std. Error	Sig. ^b
Lying & Rolling	Control (23.58)	-3.83*	0.960	0.003
	Experimental (27.41)			
Sitting	Control (27.88)	-7.22*	0.933	0.000
	Experimental (35.11)			
Crawling & Kneeling	Control (19.10)	0.09	0.846	0.920
	Experimental (19.01)			
Standing	Control (19.35)	-1.16	1.490	0.455
	Experimental (20.51)			
Walking, Running & Jogging	Control (32.04)	-2.28*	1.005	0.049
	Experimental (34.33)			
Total GMFM	Control (24.39)	-2.88*	0.822	0.007
	Experimental (27.28)			


Based on estimated marginal means

*The mean difference is significant at the .05 level.

^bAdjustment for multiple comparisons: Least Significant Difference (equivalent to no adjustments).

The pairwise comparison of the control and treatment groups shows that, there is statistically significant mean gain on GMFM dimensions; lying & rolling (MD=3.83), sitting (MD=7.22), walking, running & jogging (MD=2.28) and Total GMFM (MD=2.88). Graphical representation of the control and experimental groups pre and posttest adjusted means was presented in Figure 1.

Figure 1: Control and experimental groups' pre and post test adjusted means with standard error



L&R = Lying and rolling, C&K= Crawling and kneeling, W, R & J = Walking, Running and jumping, Total GMFM = Total gross motor function measure.

4. DISCUSSION

The analysis of the results revealed that in the case of experimental group, significant changes were seen in dimensions of GMFM 88 except in dimension C (crawling and kneeling) and D (standing) following a 16 week structured moderate physical activity programme. Statistically significant improvement was seen in dimensions A, B and E. This may probably due to the fact that exercises involving the trunk, back extensors during the conditioning activities. The lower limb muscles which are repeatedly stressed during the activities like walking, high knee action, squatting would have resulted in an increase in the strength of the muscles thus leading to an improvement in sitting.

MacPhail and Kramer, (1995) reported that significant improvements were found in dimensions D and E and Damiano and Abel, (1998) also found the results in consensus with that, the GMFM after subjects completed a strength-training program that targeted muscles of the lower limb. These sections of the GMFM measure activities such as standing alone, moving from sit to stand, walking, running, kicking, jumping, and walking up and down a step. In contrast, no change was detected in the composite GMFM score (i.e., the score from sections A–E) (Damiano, & Abel, 1998). The composite score includes items that measure lying and rolling, sitting, crawling, and kneeling, as well as those related to the activities listed above. Again, this finding is not unexpected because

training program specifically targeted lower-limb muscles; it seems reasonable to predict that increased lower limb strength would have less effect on the performance of activities such as sitting or lying and rolling than on activities such as walking, running, and jumping. It is difficult to draw from the literature conclusions about the effects of strength training on mobility in the CP population. With respect to walking speed, a positive effect after strengthening (Damiano, & Abel, 1998), whereas another study detected no change (MacPhail, & Kramer, 1998). The tailored nature of the training program may explain these disparate findings. In contrast to MacPhail and Kramer's (1998) program, which involved strengthening of the quadriceps and hamstrings regardless of each individual's assessment findings, Damiano and Abel's (1998) program targeted training of each participant's weakest lower-limb muscles. This finding suggests that strength-training programs that are tailored to individual needs may result in better functional outcomes than do less individualized programs.

It can be suggested that a training program for a minimum of 16 wks, with a frequency of three training sessions a week, may be sufficient to improve the gross motor function dimensions in CP. This finding supports the findings of Dodd, Taylor, & Damiano, (2002) and Pippenger, & Scalzitti, (2004). They conclude that there is evidence supporting the view that progressive resistance exercise can increase the ability to generate muscle force in children with CP. This conclusion was supported by another systematic review of seven studies (Darrah , Fan, Chen, Nunweiler, & Watkins, 1997).

5. CONCLUSIONS

The following conclusions were drawn on the basis of the investigation of effects of structured physical activity program for adolescents with Cerebral palsy on gross motor function measure dimensions. Based on the results obtained in the study, it can be conclude that, 16 week structured moderate physical activity program can make significant improvement in GMFM dimensions: lying and rolling, sitting and walking, running and jumping of adolescent male with cerebral palsy.

In the light of conclusions drawn, structured physical activity program must be included in the syllabi of students special education program. In future research the effect of exercise programs on the participation level in children with CP needs to be studied. Overall, only a few studies have measured the effects of an exercise program on activity in children with CP. Future studies should be conducted find effect increase in children participation in school, leisure, social and family events after undertaking an exercise program. It is surprising that no studies found examined the effects on the participation level. Especially, because participation of children with CP in everyday activities is a goal shared by

parents, service providers and organizations involved in children's rehabilitation (Law, King, King, Kertoy, Hurley, Rosenbaum, Young, & Hanna, 2006).

6. REFERENCES

- Bax, M., Goldstein, M., Rosenbaum, P., Leviton, A., Paneth, N., Dan, B., ..., & Executive Committee for the Definition of Cerebral Palsy. (2005). Proposed definition and classification of cerebral palsy, April 2005. *Developmental Medicine & Child Neurology*, 47(8), 571-576.
- Bobath, K. (1971). The normal postural reflect mechanism and its deviation in children with cerebral palsy. *Physiotherapy*, 57, 526-532.
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155-159.
- Damiano, D.L. (2006). Activity, activity, activity: rethinking our physical therapy approach to cerebral palsy. *Physical Therapy*, 86(11), 1534-1540.
- Damiano, D.L., & Abel, M.F. (1998). Functional outcomes of strength training in spastic cerebral palsy. *Archives of Physical Medicine and Rehabilitation*, 79(2), 119-125.
- Damiano, D.L., Kelly, L.E., & Vaughn, C.L. (1995). Effects of quadriceps femoris muscle strengthening on crouch gait in children with spastic diplegia. *Physical Therapy*, 75(8), 658-667.
- Darrah, J., Fan, J.S., Chen, L.C., Nunweiler, J., & Watkins, B. (1997). Review of the effects of progressive resisted muscle strengthening in children with cerebral palsy: a clinical consensus exercise. *Pediatric Physical Therapy*, 9(1), 12-17.
- Dodd, K.J., Taylor, N.F., & Damiano, D.L. (2002). A systematic review of the effectiveness of strength-training programs for people with cerebral palsy. *Archives of Physical Medicine and Rehabilitation*, 83(8), 1157-1164.
- Fowler, E.G., Ho, T.W., Nwigwe, A.I., & Dorey, F.J. (2001). The effect of quadriceps femoris muscle strengthening exercises on spasticity in children with cerebral palsy. *Physical Therapy*, 81(6), 1215-1223.
- Healy, A. (1958). Two methods of weight training for children with spastic type of cerebral palsy. *Research Quarterly*, 29, 389-395.
- Holland, L. J., & Steadward, R. D. (1990). The effects of weight training and flexibility exercising on the strength, range of motion, and spasticity/muscle tone of elite cerebral palsy athletes. In G. Doll-Tepper, C. Dahms, B. Doll & H. von Selzam (Eds.), *Adapted Physical Activity* (pp.125-129): Springer Berlin Heidelberg.
- Karande, S., & Kulkarni, M. (2008). Impact of an educational programme on parental knowledge of cerebral palsy. *Indian Journal of Paediatric*, 75(9), 901-906.
- Law, M., King, G, King, S, Kertoy, M., Hurley, P., Rosenbaum, P., Young, N., &

- Hanna, S. (2006). Patterns of participation in recreational and leisure activities among children with complex physical disabilities. *Developmental Medicine & Child Neurology*, 48(5), 337-342.
- MacPhail, H.E., & Kramer, J.F. (1995). Effect of isokinetic strength-training on functional ability and walking efficiency in adolescents with cerebral palsy. *Developmental Medicine & Child Neurology*, 37(9), 763-775.
- Paleeri, S. (2010). Educational programmes for children with cerebral palsy: parents' and teachers' dilemma. *Journal of Education and Practice*, 1(3), 9-15.
- Pippenger, W.S., & Scalzitti, D.A. (2004). What are the effects, if any, of lower-extremity strength training on gait in children with cerebral palsy. *Physical Therapy*, 84(9), 849-858.
- Russell, D.J., Rosenbaum, P.L., Wright, M., & Avery, L.M. (2013). *Gross Motor Function Measure (GMFM-66 and GMFM-88) User's Manual*, (2nd Ed.). London, UK: Mac Keith Press.
- Shepherd, R. (1995). Cerebral palsy. In R. Shepherd (ed.), *Physiotherapy in paediatrics*. (pp.110-144), Oxford, Butterworth-Heinemann.
- Stedman's Electronic Medical Dictionary*, version 4.0. (1998). Baltimore, Williams & Wilkins.
- Styer-Acevedo, J. (1999). Physical therapy for the child with cerebral palsy. *Pediatric Physical Therapy*. Philadelphia, Lippincott Williams & Wilkins.
- Verschuren, O., Ketelaar, M., Gorter, J.W., Helders, P.J., Uiterwaal, C.S., & Takken, T. (2007). Exercise training program in children and adolescents with cerebral palsy: A randomized controlled trial. *Archives of Pediatrics and Adolescent Medicine*, 161(11), 1075-1081.